



# NDST1 Rabbit pAb

|                           |  |
|---------------------------|--|
| <b>Catalog No</b>         | YP-Ab-18641  |
| <b>Isotype</b>            | IgG  |
| <b>Reactivity</b>         | Human, Mouse, Rat  |
| <b>Applications</b>       | WB   |
| <b>Gene Name</b>          | BOLA2 BOLA2A My016; BOLA2B   |
| <b>Protein Name</b>       | Bifunctional heparan sulfate N-deacetylase/N-sulfotransferase 1 (Glucosaminyl N-deacetylase/N-sulfotransferase 1) (NDST-1) (N-heparan sulfate sulfotransferase 1) (N-HSST 1) ([Heparan sulfate]-glucosamine N-sulfotransferase 1) (HSNST 1) [Includes: Heparan sulfate N-deacetylase 1 (EC 3.-.-.-); Heparan sulfate N-sulfotransferase 1 (EC 2.8.2.-)]  |
| <b>Immunogen</b>          | Synthesized peptide derived from human NDST1   |
| <b>Specificity</b>        | This antibody detects endogenous levels of NDST1 at Human, Mouse, Rat  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Source</b>             |  |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>           | WB 1:500-2000  |
| <b>Concentration</b>      | 1 mg/ml  |
| <b>Purity</b>             | ≥90%   |
| <b>Storage Stability</b>  | -20°C/1 year   |
| <b>Synonyms</b>           |  |
| <b>Observed Band</b>      | 97kD   |
| <b>Cell Pathway</b>       | [Isoform 1]: Golgi apparatus, trans-Golgi network membrane ; Single-pass type II membrane protein . Golgi apparatus, cis-Golgi network membrane ; Single-pass type II membrane protein . ; [Isoform 3]: Golgi apparatus, cis-Golgi network membrane ; Single-pass type II membrane protein .   |
| <b>Tissue Specificity</b> | Widely expressed. Expression is most abundant in heart, liver and pancreas.  |
| <b>Function</b>           | [Isoform 1]: Essential bifunctional enzyme that catalyzes both the N-deacetylation and the N-sulfation of glucosamine (GlcNAc) of the glycosaminoglycan in heparan sulfate . Modifies the GlcNAc-GlcA disaccharide repeating sugar backbone to make N-sulfated heparosan, a prerequisite substrate for later modifications in heparin biosynthesis . Plays a role in determining the extent and pattern of sulfation of heparan sulfate. Participates in biosynthesis of heparan sulfate that can ultimately serve as L-selectin ligands, thereby playing a role in inflammatory response (By similarity). Required for the exosomal release of SDCBP, CD63 and syndecan . ; [Isoform 3]: Lacks both N-deacetylase and N-sulfotransferase activities. Acts as a dominant negative on isoform 1, likely by changing the |



composition of enzyme complexes responsible for elongation and modification of heparan sulfates.

## Background

### matters needing attention

Avoid repeated freezing and thawing!

### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images