





## Fibroblast activation protein-α (FAP) Mouse mAb

Catalog No	YP-mAb-18656
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	
Protein Name	
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 411-510 of human FAP (NP_004451.2)
Specificity	
Formulation	
Source	
Purification	Affinity purification
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FAP; DPPIV; FAPA; FAPalpha; SIMP; prolyl endopeptidase FAP; Fibroblast activation protein- $\alpha$ (FAP)
Observed Band	95kDa
Cell Pathway	
Tissue Specificity	
Function	
Background	The protein encoded by this gene is a homodimeric integral membrane gelatinase belonging to the serine protease family. It is selectively expressed in reactive stromal fibroblasts of epithelial cancers, granulation tissue of healing wounds, and malignant cells of bone and soft tissue sarcomas. This protein is thought to be involved in the control of fibroblast growth or epithelial-mesenchymal interactions during development, tissue repair, and epithelial carcinogenesis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd


**€** Tel: 400-999-8863 **■** Email:3631691544@qq.com



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of various lysates, using Fibroblast activation protein-α (FAP) Mouse mAb (A23789) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.