

PADI2 Rabbit pAb

Catalog No	YP-Ab-18735
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	PAD2; PDI2; PADI2
Protein Name	Protein-arginine deiminase type-2 (PAD-H19) (Peptidylarginine deiminase II) (Protein-arginine deiminase type II)
Immunogen	Synthesized peptide derived from human PADI2
Specificity	This antibody detects endogenous levels of PADI2 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration	i ing/iii
Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 73kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 73kD Cytoplasm .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 73kD Cytoplasm . Detected in keratinocytes in epidermis (at protein level).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 73kD Cytoplasm . Detected in keratinocytes in epidermis (at protein level).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year 73kD Cytoplasm . Detected in keratinocytes in epidermis (at protein level). Catalyzes the deimination of arginine residues of proteins.



UpingBio technology Co.,Ltd



Website:	www.upingBio.com	

Pr	oducts Images