

🕓 Tel: 400-999-8863 🛛 Emall: UpingBio@163.com

Webslte: www.upingBio.com

GLRX3 Rabbit pAb

Catalog No	YP-Ab-18745
lsotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	GLRX3 PICOT TXNL2 HUSSY-22
Protein Name	Glutaredoxin-3 (PKC-interacting cousin of thioredoxin) (PICOT) (PKC-theta-interacting protein) (PKCq-interacting protein) (Thioredoxin-like protein 2)
Immunogen	Synthesized peptide derived from human GLRX3
Specificity	This antibody detects endogenous levels of GLRX3 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
	· · · · · · · · · ·
Purity	≥90%
Purity Storage Stability	
-	≥90%
Storage Stability	≥90%
Storage Stability Synonyms	≥90% -20°C/1 year
Storage Stability Synonyms Observed Band	 ≥90% -20°C/1 year 37kD Cytoplasm, cytosol . Cytoplasm, cell cortex . Cytoplasm, myofibril, sarcomere, Z line . Under the plasma membrane (By similarity). After PMA stimulation, GLRX3 and PRKCQ/PKC-theta translocate to a more extended submembrane area (By similarity). In the Z line, found associated with CSRP3 (By similarity)
Storage Stability Synonyms Observed Band Cell Pathway	 ≥90% -20°C/1 year 37kD Cytoplasm, cytosol . Cytoplasm, cell cortex . Cytoplasm, myofibril, sarcomere, Z line . Under the plasma membrane (By similarity). After PMA stimulation, GLRX3 and PRKCQ/PKC-theta translocate to a more extended submembrane area (By similarity). In the Z line, found associated with CSRP3 (By similarity) Expressed in heart, spleen, testis and, to a lower extent, in thymus and peripheral blood leukocytes. Weakly expressed in lung, placenta, colon and small intestine. Together with BOLA2, acts as a cytosolic iron-sulfur (Fe-S) cluster assembly factor that facilitates [2Fe-2S] cluster insertion into a subset of cytosolic proteins .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	 ≥90% -20°C/1 year 37kD Cytoplasm, cytosol . Cytoplasm, cell cortex . Cytoplasm, myofibril, sarcomere, Z line . Under the plasma membrane (By similarity). After PMA stimulation, GLRX3 and PRKCQ/PKC-theta translocate to a more extended submembrane area (By similarity). In the Z line, found associated with CSRP3 (By similarity) Expressed in heart, spleen, testis and, to a lower extent, in thymus and peripheral blood leukocytes. Weakly expressed in lung, placenta, colon and small intestine. Together with BOLA2, acts as a cytosolic iron-sulfur (Fe-S) cluster assembly factor that facilitates [2Fe-2S] cluster insertion into a subset of cytosolic proteins . Acts as a critical negative regulator of cardiac hypertrophy and a positive inotropic regulator (By similarity). Required for hemoglobin maturation . Does not possess any thyoredoxin activity since it lacks the conserved motif that is essential for



UpingBio technology Co.,Ltd

S Tel: 400-999-8863 Emall: UpingBio@163.com

WebsIte: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images