

S Tel: 400-999-8863 Emall: UpingBio@163.com

Webslte: www.upingBio.com

## PIR Rabbit pAb

Catalog No	YP-Ab-18762
Isotype	lgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	PIR
Protein Name	Pirin (Probable quercetin 2,3-dioxygenase PIR) (Probable quercetinase)
Immunogen	Synthesized peptide derived from human PIR
Specificity	This antibody detects endogenous levels of PIR at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 32kD
Synonyms	
Synonyms Observed Band	32kD Nucleus . Cytoplasm . Predominantly localized in dot-like subnuclear structures. Cytoplasmic localization of PIR seems to positively correlate with melanoma
Synonyms Observed Band Cell Pathway	<ul> <li>32kD</li> <li>Nucleus . Cytoplasm . Predominantly localized in dot-like subnuclear structures. Cytoplasmic localization of PIR seems to positively correlate with melanoma progression</li> <li>Highly expressed in a subset of melanomas. Detected at very low levels in most tissues (at protein level). Expressed in all tissues, with highest level of expression</li> </ul>
Synonyms Observed Band Cell Pathway Tissue Specificity	<ul> <li>32kD</li> <li>Nucleus . Cytoplasm . Predominantly localized in dot-like subnuclear structures. Cytoplasmic localization of PIR seems to positively correlate with melanoma progression</li> <li>Highly expressed in a subset of melanomas. Detected at very low levels in most tissues (at protein level). Expressed in all tissues, with highest level of expression in heart and liver.</li> <li>Transcriptional coregulator of NF-kappa-B which facilitates binding of NF-kappa-B proteins to target kappa-B genes in a redox-state-dependent manner. May be required for efficient terminal myeloid maturation of hematopoietic cells. Has</li> </ul>
Synonyms Observed Band Cell Pathway Tissue Specificity Function	<ul> <li>32kD</li> <li>Nucleus . Cytoplasm . Predominantly localized in dot-like subnuclear structures. Cytoplasmic localization of PIR seems to positively correlate with melanoma progression</li> <li>Highly expressed in a subset of melanomas. Detected at very low levels in most tissues (at protein level). Expressed in all tissues, with highest level of expression in heart and liver.</li> <li>Transcriptional coregulator of NF-kappa-B which facilitates binding of NF-kappa-B proteins to target kappa-B genes in a redox-state-dependent manner. May be required for efficient terminal myeloid maturation of hematopoietic cells. Has</li> </ul>



S Tel: 400-999-8863 Emall: UpingBio@163.com

Webslte: www.upingBio.com

## **Products Images**