







CRAT Rabbit pAb

matters needing attention	Avoid repeated freezing and thawing!
Background	
Function	Catalyzes the reversible transfer of acyl groups from carnitine to coenzyme A (CoA) and regulates the acyl-CoA/CoA ratio. Also plays a crucial role in the transport of fatty acids for beta-oxidation . Responsible for the synthesis of short-and branched-chain acylcarnitines . Active towards some branched-chain amino acid oxidation pathway (BCAAO) intermediates . Trans-2-enoyl-CoAs and 2-methylacyl-CoAs are poor substrates .
Tissue Specificity	Mostly in skeletal muscle, less in heart, liver and pancreas, only weakly detectable in brain, placenta, lung and kidney.
Cell Pathway	Endoplasmic reticulum . Peroxisome . Mitochondrion inner membrane ; Peripheral membrane protein ; Matrix side .; [Isoform 1]: Mitochondrion .; [Isoform 2]: Peroxisome .
Observed Band	69kD
Synonyms	
Storage Stability	-20°C/1 year
Purity	≥90%
Concentration	1 mg/ml
Dilution	WB 1:500-2000
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Source	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Specificity	This antibody detects endogenous levels of CRAT at Human, Mouse,Rat
Immunogen	Synthesized peptide derived from human CRAT
Protein Name	Carnitine O-acetyltransferase (Carnitine acetylase) (Carnitine acetyltransferase) (CAT) (CrAT)
Gene Name	CRAT CAT1
Applications	WB
Reactivity	Human;Rat;Mouse;
Isotype	IgG
Catalog No	YP-Ab-18763



UpingBio technology Co.,Ltd





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images