



DAAM2 Rabbit pAb

Catalog No	YP-Ab-18813
Isotype	IgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	DAAM2 KIAA0381
Protein Name	Disheveled-associated activator of morphogenesis 2
Immunogen	Synthesized peptide derived from human DAAM2
Specificity	This antibody detects endogenous levels of DAAM2 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Rabbit
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	123kD
Cell Pathway	Key regulator of the Wnt signaling pathway, which is required for various processes during development, such as dorsal patterning, determination of left/right symmetry or myelination in the central nervous system. Acts downstream of Wnt ligands and upstream of beta-catenin (CTNNB1). Required for canonical Wnt signaling pathway during patterning in the dorsal spinal cord by promoting the aggregation of Disheveled (Dvl) complexes, thereby clustering and formation of Wnt receptor signalosomes and potentiating Wnt activity. During dorsal patterning of the spinal cord, inhibits oligodendrocytes differentiation via interaction with PIP5K1A. Also regulates non-canonical Wnt signaling pathway. Acts downstream of PITX2 in the developing gut and is required for left/right asymmetry within dorsal mesentery: affects mesenchymal condensation by lengthening cadherin-based junctions through WNT5A and non-canonical Wnt signaling, inducing polarized condensation in the left dorsal mesentery necessary to initiate gut rotation. Together with DAAM1, required for myocardial maturation and sarcomere assembly. Is a regulator of actin nucleation and elongation, filopodia formation and podocyte migration .
Tissue Specificity	Cell membrane; Multi-pass membrane protein. Cell junction, gap junction.
Function	

**Background**

117kD

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Expressed in most tissues examined. Expressed in kidney glomeruli (PubMed:33232676).