



Exo1 Monoclonal Antibody

Catalog No	YP-mAb-00388
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	EXO1
Protein Name	Exonuclease 1
Immunogen	The antiserum was produced against synthesized peptide derived from human EXO1. AA range:61-110
Specificity	Exo1 Monoclonal Antibody detects endogenous levels of Exo1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EXO1; EXOI; HEX1; Exonuclease 1; hExo1; Exonuclease I; hExol
Observed Band	94kD
Cell Pathway	Nucleus . Colocalizes with PCNA to discrete nuclear foci in S-phase.
Tissue Specificity	Highly expressed in bone marrow, testis and thymus. Expressed at lower levels in colon, lymph nodes, ovary, placenta, prostate, small intestine, spleen and stomach.
Function	cofactor: Binds 2 magnesium ions per subunit. They probably participate in the reaction catalyzed by the enzyme. May bind an additional third magnesium ion after substrate binding.,developmental stage: Highly expressed in fetal liver and at lower levels in fetal brain, heart, kidney, spleen and thymus.,function: 5'→3' double-stranded DNA exonuclease which may also possess a cryptic 3'→5' double-stranded DNA exonuclease activity. Functions in DNA mismatch repair (MMR) to excise mismatch-containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. Also exhibits endonuclease activity against 5'-overhanging flap structures similar to those generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. Required for somatic hypermutation (SHM) and class switch recombination (CSR) of immunoglobulin genes. Essential for



Background

This gene encodes a protein with 5' to 3' exonuclease activity as well as an RNase H activity. It is similar to the *Saccharomyces cerevisiae* protein Exo1 which interacts with Msh2 and which is involved in mismatch repair and recombination. Alternative splicing of this gene results in three transcript variants encoding two different isoforms. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Exo1 Monoclonal Antibody