

PA26 Monoclonal Antibody

Catalog No	YP-mAb-00478
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SESN1
Protein Name	Sestrin-1
Immunogen	The antiserum was produced against synthesized peptide derived from human SESN1. AA range:271-320
Specificity	PA26 Monoclonal Antibody detects endogenous levels of PA26 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SESN1; PA26; SEST1; Sestrin-1; p53-regulated protein PA26
Observed Band	57kD
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Widely expressed.
Function	disease:Defects in SESN1 may be involved in heterotaxia. Heterotaxia is an aetiologically heterogeneous condition caused by an abnormal left-right axis formation, resulting in reversed left-right polarity of one or more organ systems.,function:Involved in the reduction of peroxiredoxins. May also be regulator of cellular growth.,induction:Isoforms T2 and isoform T3 are induced by genotoxic stress (UV, gamma-irradiation and cytotoxic drugs) in a p53-dependent manner. Isoform T1 is not induced by p53.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the sestrin family.,tissue specificity:Widely expressed.,
Background	This gene encodes a member of the sestrin family. Sestrins are induced by the p53 tumor suppressor protein and play a role in the cellular response to DNA damage and oxidative stress. The encoded protein mediates p53 inhibition of cell growth by activating AMP-activated protein kinase, which results in the inhibition of the mammalian target of rapamycin protein. The encoded protein also plays a



UpingBio technology Co.,Ltd







critical role in antioxidant defense by regenerating overoxidized peroxiredoxins, and the expression of this gene is a potential marker for exposure to radiation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PA26 Monoclonal Antibody