





## **TOB1 Monoclonal Antibody**

Catalog No	YP-mAb-00534
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	TOB1
Protein Name	Protein Tob1
Immunogen	The antiserum was produced against synthesized peptide derived from human TOB1. AA range:130-179
Specificity	TOB1 Monoclonal Antibody detects endogenous levels of TOB1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1
Storage Stability Synonyms Observed Band	-20°C/1 year  TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1  40kD  Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1  40kD  Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic  Ubiquitous.  function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1  40kD  Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic  Ubiquitous.  function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptorsPTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  TOB1; TOB; TROB1; Protein Tob1; Transducer of erbB-2 1  40kD  Cytoplasm . Nucleus . Only a small fraction localizes to the cytoplasm except in late S-phase where more than half of proteins become cytoplasmic  Ubiquitous.  function:Anti-proliferative protein that interacts with the erbB-2 receptor tyrosine kinase. May physically and/or functionally interact with protein-tyrosine kinase receptors.,PTM:Phosphorylated on Ser and Thr residues.,similarity:Belongs to the BTG family.,subunit:Interacts with CNOT7.,tissue specificity:Ubiquitous.,  This gene encodes a member of the transducer of erbB-2 /B-cell translocation gene protein family. Members of this family are anti-proliferative factors that have the potential to regulate cell growth. The encoded protein may function as a tumor suppressor. Alternate splicing results in multiple transcript variants. [provided by







**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using TOB1 Monoclonal Antibody

TOB1

-- 48

-- 117

-- 85

-- 34

(kD)