



# BCAS4 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-00682
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	BCAS4
<b>Protein Name</b>	Breast carcinoma-amplified sequence 4
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human BCAS4. AA range:31-80
<b>Specificity</b>	BCAS4 Monoclonal Antibody detects endogenous levels of BCAS4 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	BCAS4; Breast carcinoma-amplified sequence 4
<b>Observed Band</b>	23kD
<b>Cell Pathway</b>	Cytoplasm .
<b>Tissue Specificity</b>	Brain, thymus, spleen, kidney and placenta. Overexpressed in most breast cancer cell lines.
<b>Function</b>	disease:A chromosomal aberration involving BCAS4 may be a cause of breast cancer. Translocation t(17;20)(q23;q13) with BCAS3.,similarity:Belongs to the cappuccino family.,tissue specificity:Brain, thymus, spleen, kidney and placenta. Overexpressed in most breast cancer cell lines.,
<b>Background</b>	disease:A chromosomal aberration involving BCAS4 may be a cause of breast cancer. Translocation t(17;20)(q23;q13) with BCAS3.,similarity:Belongs to the cappuccino family.,tissue specificity:Brain, thymus, spleen, kidney and placenta. Overexpressed in most breast cancer cell lines.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images

