



Rabenosyn-5 Monoclonal Antibody

| | |
|---------------------------|---|
| Catalog No | YP-mAb-00718 |
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | ZFYVE20 |
| Protein Name | Rabenosyn-5 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human ZFYVE20. AA range:145-194 |
| Specificity | Rabenosyn-5 Monoclonal Antibody detects endogenous levels of Rabenosyn-5 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | ZFYVE20; Rabenosyn-5; 110 kDa protein; FYVE finger-containing Rab5 effector protein rabenosyn-5; Zinc finger FYVE domain-containing protein 20 |
| Observed Band | 89kD |
| Cell Pathway | Cell membrane; Lipid-anchor; Cytoplasmic side. Early endosome membrane ; Lipid-anchor. Enriched in endosomes that are in close proximity to clathrin-enriched regions at the cell surface. . |
| Tissue Specificity | Brain,Cervix carcinoma,Epithelium,Human salivary gland, |
| Function | function:Rab4/Rab5 effector protein acting in early endocytic membrane fusion and membrane trafficking of recycling endosomes. Required for endosome fusion either homotypically or with clathrin coated vesicles. Plays a role in the lysosomal trafficking of cathepsin D from the Golgi to lysosomes. Also promotes the recycling of transferrin directly from early endosomes to the plasma membrane. Binds phospholipid vesicles containing phosphatidylinositol 3-phosphate (PtdInsP3).,similarity:Contains 1 C2H2-type zinc finger.,similarity:Contains 1 FYVE-type zinc finger.,similarity:Contains 1 UIM (ubiquitin-interacting motif) repeat.,subunit:Interacts with EHD1, RAB4A, RAB5A, RAB22A, RAB24 and VPS45. Binds simultaneously to RAB4A and RAB5A in vitro. Interacts with RAB4A and RAB5A that has been activated by GTP binding., |

**Background**

This gene encodes a protein that belongs to the FYVE zinc finger family of proteins. The encoded protein interacts with Ras-related proteins that regulate membrane trafficking. A missense mutation in this gene is associated with a defect in the early endocytic pathway. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Rabenosyn-5 Monoclonal Antibody