



# CACNG2 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-01217
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	CACNG2
<b>Protein Name</b>	Voltage-dependent calcium channel gamma-2 subunit
<b>Immunogen</b>	Synthetic Peptide of CACNG2 AA range: 58-108
<b>Specificity</b>	The antibody detects endogenous CACNG2 protein.
<b>Formulation</b>	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Voltage-dependent calcium channel gamma-2 subunit; Neuronal voltage-gated calcium channel gamma-2 subunit; Transmembrane AMPAR regulatory protein gamma-2; TARP gamma-2
<b>Observed Band</b>	36kD
<b>Cell Pathway</b>	Membrane; Multi-pass membrane protein. Cell junction, synapse, synaptosome .
<b>Tissue Specificity</b>	Brain.
<b>Function</b>	function:Thought to stabilize the calcium channel in an inactivated (closed) state.,PTM:Phosphorylation of Thr-321 impairs interaction with DLG1 and DLG4.,similarity:Belongs to the PMP-22/EMP/MP20 family. CACNG subfamily.,subunit:The L-type calcium channel is composed of five subunits: alpha-1, alpha-2/delta, beta and gamma. Interacts with the PDZ domains of DLG4/PSD-95 and DLG1/SAP97. May interact with GOPC.,tissue specificity:Brain.,
<b>Background</b>	calcium voltage-gated channel auxiliary subunit gamma 2(CACNG2) Homo sapiens The protein encoded by this gene is a type I transmembrane AMPA receptor regulatory protein (TARP). TARPs regulate both trafficking and channel gating of the AMPA receptors. This gene is part of a functionally diverse eight-member protein subfamily of the PMP-22/EMP/MP20 family. This gene is a



susceptibility locus for schizophrenia. [provided by RefSeq, Dec 2010],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western Blot analysis of various cells using CACNG2 Monoclonal Antibody