







## CCDC102B Monoclonal Antibody

Catalog No	YP-mAb-01595
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CCDC102B
Protein Name	Coiled-coil domain-containing protein 102B
Immunogen	The antiserum was produced against synthesized peptide derived from human CCDC102B. AA range:81-130
Specificity	CCDC102B Monoclonal Antibody detects endogenous levels of CCDC102B protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CCDC102B; C18orf14; Coiled-coil domain-containing protein 102B
Observed Band	60kD
Cell Pathway	
Tissue Specificity	Lung,Salivary gland,Testis,
Function	
Background	CCDC102B (coiled-coil domain containing 102B), also known as AN, ACY1L or HsT1731, is a 513 amino acid protein that exists as three alternatively spliced isoforms. Widely expressed and found in multiple CNV (copy-number variant) regions, CCDC102B contains the deletion breakpoint of a maternally inherited deletion, which is 2.7 Mb in size, and maps to human chromosome 18q22.1. CCDC102B may play a role in the pathogenesis of diaphragmatic hernia, microphthalmia, colorectal carcinoma and schizophrenia. Encoding over 300 genes, chromosome 18 contains about 76 million bases. Translocation between chromosomes 18 and 14 is the most common translocation in cancers and occurs in follicular lymphomas. Niemann-Pick disease, hereditary hemorrhagic telangiectasia and erythropoietic protoporphyria are associated with chromosome 18.



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using CCDC102B Monoclonal Antibody