

CUX1 Monoclonal Antibody

Catalog No	YP-mAb-01603
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	CUX1
Protein Name	Homeobox protein cut-like 1/Protein CASP
Immunogen	The antiserum was produced against synthesized peptide derived from human CUTL1. AA range:10-59
Specificity	CUX1 Monoclonal Antibody detects endogenous levels of CUX1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CUX1; CUTL1; Homeobox protein cut-like 1; CCAAT displacement protein; CDP; Homeobox protein cux-1; CUX1; CUTL1; Protein CASP
Observed Band	164kD
Cell Pathway	Nucleus .
Tissue Specificity	Colon,Duodenum,Liver,Umbilical vein,
Function	alternative products:Additional isoforms seem to exist,function:May be involved in intra-Golgi retrograde transport.,function:Probably has a broad role in mammalian development as a repressor of developmentally regulated gene expression. May act by preventing binding of positively-activing CCAAT factors to promoters. Component of nf-munr repressor; binds to the matrix attachment regions (MARs) (5' and 3') of the immunoglobulin heavy chain enhancer. Represses T-cell receptor (TCR) beta enhancer function by binding to MARbeta, an ATC-rich DNA sequence located upstream of the TCR beta enhancer.,miscellaneous:Asn-1290 may participate in regulating DNA-binding activity by promoting homo- and heterodimerization.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the CASP family.,similarity:Belongs to the CUT homeobox family.,similarity:Contains 1 homeobox DNA-bin



UpingBio technology Co.,Ltd

▼ Tel: 400-999-8863
Emall: UpingBio@163.com



Background	The protein encoded by this gene is a member of the homeodomain family of DNA binding proteins. It may regulate gene expression, morphogenesis, and differentiation and it may also play a role in the cell cycle progession. Several alternatively spliced transcript variants encoding different isoforms have been identified.[provided by RefSeq, Feb 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

