



CtBP2 Monoclonal Antibody

Catalog No	YP-mAb-01634
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CTBP2
Protein Name	C-terminal-binding protein 2
Immunogen	The antiserum was produced against synthesized peptide derived from human CtBP2. AA range:396-445
Specificity	CtBP2 Monoclonal Antibody detects endogenous levels of CtBP2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CTBP2; C-terminal-binding protein 2; CtBP2
Observed Band	49kD
Cell Pathway	Nucleus . Cell junction, synapse .
Tissue Specificity	Ubiquitous. Highest levels in heart, skeletal muscle, and pancreas.
Function	function:Corepressor targeting diverse transcription regulators. Isoform 2 probably acts as a scaffold for specialized synapses.,PTM:Isoform 2 is phosphorylated upon DNA damage, probably by ATM or ATR at Thr-179; Ser-181 and Ser-185. Phosphorylation by HIPK2 on Ser-428 induces proteasomal degradation.,similarity:Belongs to the D-isomer specific 2-hydroxyacid dehydrogenase family.,subunit:Interacts with the C-terminus of adenovirus E1A protein. Can form homodimers or heterodimers of CTBP1 and CTBP2. Interacts with HIPK2 (By similarity). Interacts with PNN, NRIP1 and WIZ.,tissue specificity:Ubiquitous. Highest levels in heart, skeletal muscle, and pancreas.,
Background	This gene produces alternative transcripts encoding two distinct proteins. One protein is a transcriptional repressor, while the other isoform is a major component of specialized synapses known as synaptic ribbons. Both proteins contain a NAD ⁺ binding domain similar to NAD ⁺ -dependent 2-hydroxyacid dehydrogenases. A portion of the 3' untranslated region was used to map this gene to



chromosome 21q21.3; however, it was noted that similar loci elsewhere in the genome are likely. Blast analysis shows that this gene is present on chromosome 10. Several transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Feb 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CtBP2 Monoclonal Antibody