

## DNA pol a Monoclonal Antibody

Catalog No	YP-mAb-01654
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	POLA1
Protein Name	DNA polymerase alpha catalytic subunit
Immunogen	The antiserum was produced against synthesized peptide derived from human DNA Polymerase alpha. AA range:81-130
Specificity	DNA pol $^{ \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
	3
Purity	≥90%
Purity Storage Stability	
	≥90%
Storage Stability	≥90%  -20°C/1 year  POLA1; POLA; DNA polymerase alpha catalytic subunit; DNA polymerase alpha
Storage Stability Synonyms	≥90%  -20°C/1 year  POLA1; POLA; DNA polymerase alpha catalytic subunit; DNA polymerase alpha catalytic subunit p180
Storage Stability Synonyms Observed Band	≥90%  -20°C/1 year  POLA1; POLA; DNA polymerase alpha catalytic subunit; DNA polymerase alpha catalytic subunit p180  165kD  Nucleus . Cytoplasm, cytosol . In the cytosol, colocalizes with RNA:DNA hybrids
Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  POLA1; POLA; DNA polymerase alpha catalytic subunit; DNA polymerase alpha catalytic subunit p180  165kD  Nucleus . Cytoplasm, cytosol . In the cytosol, colocalizes with RNA:DNA hybrids
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  POLA1; POLA; DNA polymerase alpha catalytic subunit; DNA polymerase alpha catalytic subunit p180  165kD  Nucleus . Cytoplasm, cytosol . In the cytosol, colocalizes with RNA:DNA hybrids with a speckled pattern  catalytic activity:Deoxynucleoside triphosphate + DNA(n) = diphosphate + DNA(n+1).,function:Polymerase alpha in a complex with DNA primase is a replicative polymerase.,miscellaneous:In eukaryotes there are five DNA polymerases: alpha, beta, gamma, delta, and epsilon which are responsible for different reactions of DNA synthesis.,PTM:A 165 kDa form is probably produced by proteolytic cleavage at Lys-124.,similarity:Belongs to the DNA polymerase type-B family.,subunit:Interacts with SV40 Large T antigen; this interaction allows



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using DNA pol α Monoclonal Antibody