



DNA pol β Monoclonal Antibody

Catalog No	YP-mAb-01655
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	POLB
Protein Name	DNA polymerase beta
Immunogen	The antiserum was produced against synthesized peptide derived from human DNA Polymerase beta. AA range:286-335
Specificity	DNA pol β Monoclonal Antibody detects endogenous levels of DNA pol β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	POLB; DNA polymerase beta
Observed Band	38kD
Cell Pathway	Nucleus. Cytoplasm. Cytoplasmic in normal conditions. Translocates to the nucleus following DNA damage.
Tissue Specificity	Skin,Testis,
Function	catalytic activity:Deoxynucleoside triphosphate + DNA(n) = diphosphate + DNA(n+1).,cofactor: Binds 2 magnesium ions per subunit.,domain:Residues 239-252 form a flexible loop which appears to affect the polymerase fidelity.,function:Repair polymerase. Conducts "gap-filling" DNA synthesis in a stepwise distributive fashion rather than in a processive fashion as for other DNA polymerases. Has a 5'-deoxyribose-5-phosphate lyase (dRP lyase) activity.,PTM:Methylation by PRMT6 stimulates the polymerase activity by enhancing DNA binding and processivity.,similarity:Belongs to the DNA polymerase type-X family.,subunit:Monomer.,
Background	The protein encoded by this gene is a DNA polymerase involved in base excision and repair, also called gap-filling DNA synthesis. The encoded protein, acting as a monomer, is normally found in the cytoplasm, but it translocates to the nucleus upon DNA damage. Several transcript variants of this gene exist, but the full-length nature of only one has been described to date. [provided by RefSeq,



Sep 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DNA pol β Monoclonal Antibody