

EFP Monoclonal Antibody

Catalog No	YP-mAb-01682
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TRIM25
Protein Name	E3 ubiquitin/ISG15 ligase TRIM25
Immunogen	The antiserum was produced against synthesized peptide derived from human ZNF147. AA range:201-250
Specificity	EFP Monoclonal Antibody detects endogenous levels of EFP protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TRIM25; EFP; RNF147; ZNF147; E3 ubiquitin/ISG15 ligase TRIM25; Estrogen-responsive finger protein; RING finger protein 147; Tripartite motif-containing protein 25; Ubiquitin/ISG15-conjugating enzyme TRIM25; Zinc finger protein 147
Observed Band	71kD
Cell Pathway	Cytoplasm . Cytoplasm, Stress granule . Nucleus .
Tissue Specificity	Expressed in breast tumors (at protein level). Ubiquitous.
Function	function:Mediates estrogen action in various target organs.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,tissue specificity:Ubiquitous.,
Background	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to the cytoplasm. The presence of potential DNA-binding and dimerization-transactivation domains suggests that this protein may act as a transcription factor, similar to several other members of the TRIM family. Expression of the gene is upregulated in response to estrogen, and it is thought to mediate estrogen actions in breast cancer as a primary response gene. [provided



UpingBio technology Co.,Ltd







by RefSeq, Jul 2008],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

