



GCN5 Monoclonal Antibody

Catalog No	YP-mAb-01748
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	KAT2A
Protein Name	Histone acetyltransferase KAT2A
Immunogen	The antiserum was produced against synthesized peptide derived from human GCN5L2. AA range:691-740
Specificity	GCN5 Monoclonal Antibody detects endogenous levels of GCN5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KAT2A; GCN5; GCN5L2; HGCN5; Histone acetyltransferase KAT2A; General control of amino acid synthesis protein 5-like 2; Histone acetyltransferase GCN5; HsGCN5; Lysine acetyltransferase 2A; STAF97
Observed Band	100kD
Cell Pathway	Nucleus . Chromosome . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Mainly localizes to the nucleus (PubMed:27796307). Also localizes to centrosomes in late G1 and around the G1/S transition, coinciding with the onset of centriole formation (PubMed:27796307). .
Tissue Specificity	Expressed in all tissues tested.
Function	somitogenesis, regionalization, chromatin organization, chromatin remodeling, transcription, transcription, DNA-dependent, regulation of transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter,transcription from RNA polymerase II promoter, protein amino acid acetylation, pattern specification process,embryonic development ending in birth or egg hatching, anterior/posterior pattern formation, chromatin modification,covalent chromatin modification, histone modification, histone acetylation, histone deubiquitination, protein deubiquitination, RNA biosynthetic process, segmentation, chordate embryonic development, protein amino acid acylation, histone H3 acetylation, regulation of transcription, regulation of RNA metabolic process, chromosome organization, protein modification by small



protein removal, protein modification by small protein conjugation or rem

Background

KAT2A, or GCN5, is a histone acetyltransferase (HAT) that functions primarily as a transcriptional activator. It also functions as a repressor of NF-kappa-B (see MIM 164011) by promoting ubiquitination of the NF-kappa-B subunit RELA (MIM 164014) in a HAT-independent manner (Mao et al., 2009 [PubMed 19339690]).[supplied by OMIM, Sep 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GCN5 Monoclonal Antibody