



IRF-2 Monoclonal Antibody

Catalog No	YP-mAb-01825
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	IRF2
Protein Name	Interferon regulatory factor 2
Immunogen	The antiserum was produced against synthesized peptide derived from human IRF2. AA range:101-150
Specificity	IRF-2 Monoclonal Antibody detects endogenous levels of IRF-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IRF2; Interferon regulatory factor 2; IRF-2
Observed Band	47kD
Cell Pathway	Nucleus.
Tissue Specificity	Expressed throughout the epithelium of the colon. Also expressed in lamina propria.
Function	function:Specifically binds to the upstream regulatory region of type I IFN and IFN-inducible MHC class I genes (the interferon consensus sequence (ICS)) and represses those genes. Also acts as an activator for several genes including H4 and IL7. Constitutively binds to the ISRE promoter to activate IL7. Involved in cell cycle regulation through binding the site II (HiNF-M) promoter region of H4 and activating transcription during cell growth. Antagonizes IRF1 transcriptional activation.,induction:By viruses and IFN.,PTM:Acetylated by CBP/ p300 during cell-growth. Acetylation on Lys-75 is required for stimulation of H4 promoter activity.,PTM:The major sites of sumoylation are Lys-137 and Lys-293. Sumoylation by SUMO1 increases its transcriptional repressor activity on IRF1 and diminishes its ability to activate ISRE and H4 promoter.,similarity:Belongs to the IRF family.,similarity:Contai
Background	IRF2 encodes interferon regulatory factor 2, a member of the interferon regulatory transcription factor (IRF) family. IRF2 competitively inhibits the



IRF1-mediated transcriptional activation of interferons alpha and beta, and presumably other genes that employ IRF1 for transcription activation. However, IRF2 also functions as a transcriptional activator of histone H4. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

