



# Med8 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-01863
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	MED8
<b>Protein Name</b>	Mediator of RNA polymerase II transcription subunit 8
<b>Immunogen</b>	Synthesized peptide derived from Med8 . at AA range: 70-150
<b>Specificity</b>	Med8 Monoclonal Antibody detects endogenous levels of Med8 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	MED8; Mediator of RNA polymerase II transcription subunit 8; Activator-recruited cofactor 32 kDa component; ARC32; Mediator complex subunit 8
<b>Observed Band</b>	29kD
<b>Cell Pathway</b>	Nucleus .
<b>Tissue Specificity</b>	Cervix carcinoma,Melanoma,Testis,
<b>Function</b>	domain:The elongin BC complex binding domain is also known as BC-box with the consensus [APST]-L-x(3)-C-x(3)-[AIV]..function:Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May play a role as a target recruitment subunit in E3 ubiquitin-protein ligase complexes and thus in ubiquitination and subsequent proteasomal degradation of target proteins..pathway:Protein modification; protein ubiquitination..similarity:Belongs to the Mediator
<b>Background</b>	This gene encodes a protein component of the mediator complex, which aids in transcriptional activation through interaction with RNA polymerase II and



gene-specific transcription factors. The encoded protein may also function in ubiquitin ligation and protein degradation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2013],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using Med8 Monoclonal Antibody