



mTERF Monoclonal Antibody

Catalog No	YP-mAb-01882
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	MTERF
Protein Name	Transcription termination factor mitochondrial
Immunogen	The antiserum was produced against synthesized peptide derived from human MTERF. AA range:267-316
Specificity	mTERF Monoclonal Antibody detects endogenous levels of mTERF protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MTERF; Transcription termination factor; mitochondrial; Mitochondrial transcription termination factor 1; mTERF
Observed Band	46kD
Cell Pathway	Mitochondrion.
Tissue Specificity	Brain,Colon,Placenta,Uterus,
Function	domain:Composed of three leucine zippers, one of which is bipartite, and two widely spaced basic domains. There is evidence that the leucine zippers form and intramolecular three-stranded coiled-coil that brings the basic domains together to form a DNA-binding motif.,function:Transcription termination factor. Binds to a 28 bp region within the tRNA(Leu(uur)) gene at a position immediately adjacent to and downstream of the 16S rRNA gene, this region comprises a tridecamer sequence critical for directing accurate termination. Probably requires one or more components for termination activity.,PTM:Phosphoprotein with mostly four phosphate groups. While the DNA-binding activity is unaffected by the phosphorylation state, only the phosphorylated form of the protein is active for termination activity. Functioning seems to be regulated by phosphorylation.,similarity:Belongs to the mTERF family.,



Background

This gene encodes a mitochondrial transcription termination factor. This protein participates in attenuating transcription from the mitochondrial genome; this attenuation allows higher levels of expression of 16S ribosomal RNA relative to the tRNA gene downstream. The product of this gene has three leucine zipper motifs bracketed by two basic domains that are all required for DNA binding. There is evidence that, for this protein, the zippers participate in intramolecular interactions that establish the three-dimensional structure required for DNA binding. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

