



Nrl Monoclonal Antibody

| Catalog No | YP-mAb-01920 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | NRL |
| Protein Name | Neural retina-specific leucine zipper protein |
| Immunogen | The antiserum was produced against synthesized peptide derived from human NRL. AA range:19-68 |
| Specificity | Nrl Monoclonal Antibody detects endogenous levels of Nrl protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | NRL; D14S46E; Neural retina-specific leucine zipper protein; NRL |
| Observed Band | 25kD |
| Cell Pathway | Cytoplasm . Nucleus . |
| Tissue Specificity | Expressed in the brain and the retina (PubMed:11477108). Expressed strongly in rod and cone cells (at protein level) (PubMed:11477108). |
| Function | disease:Defects in NRL are the cause of retinitis pigmentosa type 27 (RP27) [MIM:162080]. RP leads to degeneration of retinal photoreceptor cells. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well. RP27 inheritance is autosomal dominant.,function:Transcription factor which regulates the expression of several rod-specific genes, in cluding RHO and PDE6B.,online information:Retina International's Scientific Newsletter,similarity:Belongs to the bZIP family.,similarity:Contains 1 bZIP domain.,subunit:Interacts with FIZ1. This interaction represses transactivation.,tissue specificity:Neural retina., |
| Background | This gene encodes a basic motif-leucine zipper transcription factor of the Maf subfamily. The encoded protein is conserved among vertebrates and is a critical intrinsic regulator of photoceptor development and function. Mutations in this gene |

have been associated with retinitis pigmentosa and retinal degenerative diseases.



UpingBio technology Co.,Ltd








[provided by RefSeq, Jul 2008],

| matters needing attention | Avoid repeated freezing and thawing! |
|---------------------------|--|
| Usage suggestions | This product can be used in immunological reaction |

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Nrl Monoclonal Antibody