

ODC Monoclonal Antibody

Catalog No YP-mAb-01927 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name SLC25A21 Protein Name Mitochondrial 2-oxodicarboxylate carrier Immunogen The antiserum was produced against synthesized peptide derived from human SLC25A21. AA range:134-183 Specificity ODC Monoclonal Antibody detects endogenous levels of ODC protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transport Sc 5-C7 oxodicarboxylates across the i		
Reactivity Human; Mouse; Rat Applications WB Gene Name SLC25A21 Protein Name Mitochondrial 2-oxodicarboxylate carrier Immunogen The antiserum was produced against synthesized peptide derived from human SLC25A21. AA range: 134-183 Specificity ODC Monoclonal Antibody detects endogenous levels of ODC protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function: Transports CS-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxogilutarate, adipate, glutarate, and to a lesser extent, pinelate, 2-oxopimelate, 2-axinoadipate, oxaloacetate, and citrate, similarity, Belongs to the mitochondrial carriers that transport CS-C7 oxodicarboxylates across inner mitochondrial carriers	Catalog No	YP-mAb-01927
Applications Gene Name SLC25A21 Protein Name Mitochondrial 2-oxodicarboxylate carrier Immunogen The antiserum was produced against synthesized peptide derived from human SLC25A21. AA range:134-183 Specificity ODC Monoclonal Antibody detects endogenous levels of ODC protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function: Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and citrate, similarity Belongs to the mitochondrial carrier family, similarity: Contains 3 Solcar repeats, tissue specificity: Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carrierers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria. Convactionetic sonverted into	Isotype	IgG
Gene Name SLC25A21 Protein Name Mitochondrial 2-oxodicarboxylate carrier Immunogen The antiserum was produced against synthesized peptide derived from human SLC25A21. AA range:134-183 Specificity ODC Monoclonal Antibody detects endogenous levels of ODC protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and citrate, similarity. Belongs to the mitochondrial carrier family, similarity. Contains 3 Solcar repeats, tissue specificity:Ubiquitous, Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria! scorovered into	Reactivity	Human;Mouse;Rat
Protein Name Mitochondrial 2-oxodicarboxylate carrier	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human SLC25A21. AA range:134-183 Specificity ODC Monoclonal Antibody detects endogenous levels of ODC protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate, similarity. Belongs to the mitochondrial carrier family, similarity. Contains 3 Solcar repeats, tissue specificity:Ubiquitous. Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within interhodnerical transported into	Gene Name	SLC25A21
SLC25A21. AA range:134-183 Specificity ODC Monoclonal Antibody detects endogenous levels of ODC protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-oxoglutarate, adipate, glutarate, and citrate, similarity:Belongs to the mitochondrial carrier family, similarity:Contains 3 Solcar repeats, tissue specificity: Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Protein Name	Mitochondrial 2-oxodicarboxylate carrier
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate, similarity:Belongs to the mitochondrial carrier familysimilarity:Contains 3 Solcar repeats, tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and citrate, similarity:Belongs to the mitochondrial carrier family.,similarity:Contains 3 Solcar repeats.,tissue specificity:Ubiquitous. Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria 2 coxoadipate is converted into	Specificity	ODC Monoclonal Antibody detects endogenous levels of ODC protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity: Belongs to the mitochondrial carrier family., similarity:Contains 3 Solcar repeats., tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial arrieriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria; a converted into	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and citrate, similarity:Belongs to the mitochondrial carrier family, similarity:Contains 3 Solcar repeats, tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria. 2-oxoadipate is converted into	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citratesimilarity:Belongs to the mitochondrial carrier familysimilarity:Contains 3 Solcar repeats, tissue specificity:Ubiquitous. Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity:Belongs to the mitochondrial carrier family., similarity:Contains 3 Solcar repeats., tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Dilution	WB 1:500-1:2000
Storage Stability -20°C/1 year Sunonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band Gell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function: Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity: Belongs to the mitochondrial carrier family., similarity: Contains 3 Solcar repeats., tissue specificity: Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Concentration	1 mg/ml
Synonyms SLC25A21; ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21 Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity:Belongs to the mitochondrial carrier family., similarity:Contains 3 Solcar repeats., tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Purity	≥90%
Observed Band 36kD Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Expressed in placenta, gall bladder and colon. Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity:Belongs to the mitochondrial carrier family., similarity:Contains 3 Solcar repeats., tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Storage Stability	-20°C/1 year
Cell Pathway Mitochondrion inner membrane; Multi-pass membrane protein. Expressed in placenta, gall bladder and colon. function: Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity: Belongs to the mitochondrial carrier family., similarity: Contains 3 Solcar repeats., tissue specificity: Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Synonyms	•
Tissue Specificity Expressed in placenta, gall bladder and colon. function: Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity: Belongs to the mitochondrial carrier family., similarity: Contains 3 Solcar repeats., tissue specificity: Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Observed Band	36kD
Function function:Transports C5-C7 oxodicarboxylates across the inner membranes of mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate.,similarity:Belongs to the mitochondrial carrier family.,similarity:Contains 3 Solcar repeats.,tissue specificity:Ubiquitous., SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Cell Pathway	Mitochondrion inner membrane; Multi-pass membrane protein.
mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity:Belongs to the mitochondrial carrier family., similarity:Contains 3 Solcar repeats., tissue specificity:Ubiquitous., Background SLC25A21 is a homolog of the S. cerevisiae ODC proteins, mitochondrial carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Tissue Specificity	Expressed in placenta, gall bladder and colon.
carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into	Function	mitochondria. Can transport 2-oxoadipate, 2-oxoglutarate, adipate, glutarate, and to a lesser extent, pimelate, 2-oxopimelate, 2-aminoadipate, oxaloacetate, and citrate., similarity:Belongs to the mitochondrial carrier family., similarity:Contains 3
	Background	carriers that transport C5-C7 oxodicarboxylates across inner mitochondrial membranes. One of the species transported by ODC is 2-oxoadipate, a common intermediate in the catabolism of lysine, tryptophan, and hydroxylysine in mammals. Within mitochondria, 2-oxoadipate is converted into



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ODC Monoclonal Antibody