



# RFX3 Monoclonal Antibody


<b>Catalog No</b>	YP-mAb-01980
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	RFX3
<b>Protein Name</b>	Transcription factor RFX3
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human RFX3. AA range:641-690
<b>Specificity</b>	RFX3 Monoclonal Antibody detects endogenous levels of RFX3 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	RFX3; Transcription factor RFX3; Regulatory factor X 3
<b>Observed Band</b>	84kD
<b>Cell Pathway</b>	Nucleus .
<b>Tissue Specificity</b>	Muscle,Testis,
<b>Function</b>	function:Acts as a transcription factor. Represses transcription of MAP1A in non-neuronal cells but not in neuronal cells.,similarity:Belongs to the RFX family.,
<b>Background</b>	This gene is a member of the regulatory factor X gene family, which encodes transcription factors that contain a highly-conserved winged helix DNA binding domain. The protein encoded by this gene is structurally related to regulatory factors X1, X2, X4, and X5. It is a transcriptional activator that can bind DNA as a monomer or as a heterodimer with other RFX family members. Multiple transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Aug 2013],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western Blot analysis of various cells using RFX3 Monoclonal Antibody