

## RPA40 Monoclonal Antibody

Catalog No	YP-mAb-01993
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	POLR1C
Protein Name	DNA-directed RNA polymerases I and III subunit RPAC1
Immunogen	The antiserum was produced against synthesized peptide derived from human POLR1C. AA range:101-150
Specificity	RPA40 Monoclonal Antibody detects endogenous levels of RPA40 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	POLR1C; POLR1E; DNA-directed RNA polymerases I and III subunit RPAC1; DNA-directed RNA polymerase I subunit C; RNA polymerases I and III subunit AC1; AC40; DNA-directed RNA polymerases I and III 40 kDa polypeptide; RPA40; RPA39; RPC40
Observed Band	39kD
Cell Pathway	Nucleus .
Tissue Specificity	Brain,Ovarian carcinoma,Placenta,Umbilical cord blood,
Function	function:DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Common component of RNA polymerases I and III which synthesize ribosomal RNA precursors and small RNAs, such as 5S rRNA and tRNAs, respectively. RPAC1 is part of the Pol core element with the central large cleft and probably a clamp element that moves to open and close the cleft.,similarity:Belongs to the archaeal rpoD/eukaryotic RPB3 RNA polymerase subunit family.,subunit:Component of the RNA polymerase I (Pol I) and RNA polymerase III (Pol III) complexes consisting of at least 13 and 17 subunits, respectively.,



## UpingBio technology Co.,Ltd







Background	The protein encoded by this gene is a subunit of both RNA polymerase I and RNA polymerase III complexes. The encoded protein is part of the Pol core element. Mutations in this gene have been associated with Treacher Collins syndrome (TCS) and hypomyelinating leukodystrophy 11. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



