



HNF-1 β Monoclonal Antibody

Catalog No	YP-mAb-02222
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HNF1B
Protein Name	Hepatocyte nuclear factor 1-beta
Immunogen	Synthesized peptide derived from the N-terminal region of human HNF-1 β .
Specificity	HNF-1 β Monoclonal Antibody detects endogenous levels of HNF-1 β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	HNF1B; TCF2; Hepatocyte nuclear factor 1-beta; HNF-1-beta; HNF-1B; Homeoprotein LFB3; Transcription factor 2; TCF-2; Variant hepatic nuclear factor 1; vHNF1
Observed Band	60kD
Cell Pathway	Nucleus.
Tissue Specificity	Colon,Liver,Thalamus,
Function	disease:A genetic variation in HNF1B is associated with susceptibility to hereditary prostate cancer type 11 (HPC11) [MIM:611955].,disease:Defects in HNF1B are a cause of Muellerian aplasia [MIM:158330]. In a Norwegian family with a novel syndrome of mild diabetes and severe non-diabetic renal disease, Muellerian aplasia expressed as vaginal aplasia and rudimentary uterus, were found in 2 females. These findings suggest that a broader spectrum of clinical symptoms may be associated with defects in HNF1B than previously recognized.,disease:Defects in HNF1B are the cause of maturity-onset diabetes of the young type 5 (MODY5) [MIM:604284]. MODY [MIM:606391] is a form of diabetes mellitus characterized by an autosomal dominant mode of inheritance, age of onset of 25 years or younger and a primary defect in insulin secretion.,disease:Defects in HNF1B are the cause of renal cysts and diabetes



Background

This gene encodes a member of the homeodomain-containing superfamily of transcription factors. The protein binds to DNA as either a homodimer, or a heterodimer with the related protein hepatocyte nuclear factor 1-alpha. The gene has been shown to function in nephron development, and regulates development of the embryonic pancreas. Mutations in this gene result in renal cysts and diabetes syndrome and noninsulin-dependent diabetes mellitus, and expression of this gene is altered in some types of cancer. Multiple transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Sep 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using HNF-1 β Monoclonal Antibody