



# Topo III $\alpha$ Monoclonal Antibody

Catalog No	YP-mAb-02256
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TOP3A
Protein Name	DNA topoisomerase 3-alpha
Immunogen	Synthesized peptide derived from Topo III $\alpha$ . at AA range: 350-430
Specificity	Topo III $\alpha$ Monoclonal Antibody detects endogenous levels of Topo III $\alpha$ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	TOP3A; TOP3; DNA topoisomerase 3-alpha; DNA topoisomerase III alpha
Observed Band	115kD
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	High expression is found in testis, heart, skeletal muscle and pancreas.
Function	catalytic activity:ATP-independent breakage of single-stranded DNA, followed by passage and rejoining.,function:Reduces the number of supercoils in a highly negatively supercoiled DNA. Essential component of the RMI complex, a complex that plays an important role in the processing of homologous recombination intermediates to limit DNA crossover formation in cells.,similarity:Belongs to the prokaryotic type I/III topoisomerase family.,subunit:Directly interacts with BLM and RMI1. Component of the RMI complex, containing at least TOP3A, RMI1 and RMI2. The RMI complex interacts with BLM.,tissue specificity:High expression is found in testis, heart, skeletal muscle and pancreas.,
Background	This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This enzyme catalyzes the transient breaking and rejoining of a single strand of DNA which allows the strands to pass through one another, thus reducing the number of supercoils and altering the topology of DNA. This enzyme forms a complex with BLM which functions in the regulation of recombination in somatic cells. Alternative splicing



results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Mar 2016],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using Topo III  $\alpha$  Monoclonal Antibody