



CHSY2 Monoclonal Antibody

Catalog No	YP-mAb-02536
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CHPF
Protein Name	Chondroitin sulfate synthase 2
Immunogen	The antiserum was produced against synthesized peptide derived from human CHSY2. AA range:631-680
Specificity	CHSY2 Monoclonal Antibody detects endogenous levels of CHSY2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CHPF; CSS2; Chondroitin sulfate synthase 2; Chondroitin glucuronyltransferase 2; Chondroitin-polymerizing factor; ChPF; Glucuronosyl-N-acetylgalactosaminyl-proteoglycan 4-beta-N-acetylgalactosaminyltransferase II; N-acetylgalactosaminyl-pro
Observed Band	85kD
Cell Pathway	[Isoform 1]: Golgi apparatus, Golgi stack membrane ; Single-pass type II membrane protein . Cytoplasm, cytosol .; [Isoform 3]: Cytoplasm, cytosol . Mitochondrion .; [Isoform 2]: Mitochondrion matrix .
Tissue Specificity	Ubiquitous. Highly expressed in pancreas, ovary, brain, heart, skeletal muscle, colon, kidney, liver, stomach, spleen and placenta. ; [Isoform 2]: Expressed in brain, spleen, ovary, testis, lung and peripheral mononuclear cells. ; [Isoform 3]: Also ubiquitous.
Function	catalytic activity:UDP-alpha-D-glucuronate + N-acetyl-beta-D-galactosaminyl-(1->4)-beta-D-glucuronosyl-proteoglycan = UDP + beta-D-glucuronosyl-(1->3)-N-acetyl-beta-D-galactosaminyl-(1->4)-beta-D-glucuronosyl-proteoglycan.,catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronosyl-(1->3)-N-acetyl-beta-D-galactosaminyl-proteoglycan = UDP + N-acetyl-beta-D-galactosaminyl-(1->4)-beta-D-glucuronosyl-(1->3)-N-acetyl-beta-



D-galactosaminyl-proteoglycan.,cofactor:Divalent cations. Highest activities are measured with manganese. Can also utilize cobalt.,function:Has both beta-1,3-glucuronic acid and beta-1,4-N-acetylgalactosamine transferase activity. Transfers glucuronic acid (GlcUA) from UDP-GlcUA and N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of the elongating chondroitin polymer.,online information:GlycoGene database,PTM:Phosphorylated upon DNA damage, p

Background

catalytic activity:UDP-alpha-D-glucuronate + N-acetyl-beta-D-galactosaminyl-(1->4)-beta-D-glucuronosyl-proteoglycan = UDP + beta-D-glucuronosyl-(1->3)-N-acetyl-beta-D-galactosaminyl-(1->4)-beta-D-glucuronosyl-proteoglycan.,catalytic activity:UDP-N-acetyl-D-galactosamine + beta-D-glucuronosyl-(1->3)-N-acetyl-beta-D-galactosaminyl-proteoglycan = UDP + N-acetyl-beta-D-galactosaminyl-(1->4)-beta-D-glucuronosyl-(1->3)-N-acetyl-beta-D-galactosaminyl-proteoglycan.,cofactor:Divalent cations. Highest activities are measured with manganese. Can also utilize cobalt.,function:Has both beta-1,3-glucuronic acid and beta-1,4-N-acetylgalactosamine transferase activity. Transfers glucuronic acid (GlcUA) from UDP-GlcUA and N-acetylgalactosamine (GalNAc) from UDP-GalNAc to the non-reducing end of the elongating chondroitin polymer.,online information:GlycoGene database,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the chondroitin N-acetylgalactosaminyltransferase family.,subunit:Binds CHSY1.,tissue specificity:Ubiquitous. Highly expressed in pancreas, ovary, brain, heart, skeletal muscle, colon, kidney, liver, stomach, small intestine and placenta.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

