



# DECR2 Monoclonal Antibody

Catalog No	YP-mAb-02613
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	DECR2
Protein Name	Peroxisomal 2,4-dienoyl-CoA reductase
Immunogen	The antiserum was produced against synthesized peptide derived from human DECR2. AA range:217-266
Specificity	DECR2 Monoclonal Antibody detects endogenous levels of DECR2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DECR2; PDCR; Peroxisomal 2; 4-dienoyl-CoA reductase; pDCR; 2,4-dienoyl-CoA reductase 2
Observed Band	36kD
Cell Pathway	Peroxisome .
Tissue Specificity	Brain,Epithelium,Lung,Testis,
Function	catalytic activity:Trans-2,3-didehydroacyl-CoA + NADP(+) = trans,trans-2,3,4,5-tetradehydroacyl-CoA + NADPH.,function:Auxiliary enzyme of beta-oxidation. Participates in the degradation of unsaturated fatty enoyl-CoA esters having double bonds in both even- and odd-numbered positions in peroxisome. Catalyzes the NADP-dependent reduction of 2,4-dienoyl-CoA to yield trans-3-enoyl-CoA. Has activity towards short and medium chain 2,4-dienoyl-CoAs, but also towards 2,4,7,10,13,16,19-docosaheptaenoyl-CoA, suggesting that it does not constitute a rate limiting step in the peroxisomal degradation of docosahexaenoic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. 2,4-dienoyl-CoA reductase subfamily.,
Background	catalytic activity:Trans-2,3-didehydroacyl-CoA + NADP(+) = trans,trans-2,3,4,5-tetradehydroacyl-CoA + NADPH.,function:Auxiliary enzyme of beta-oxidation. Participates in the degradation of unsaturated fatty enoyl-CoA



esters having double bonds in both even- and odd-numbered positions in peroxisome. Catalyzes the NADP-dependent reduction of 2,4-dienoyl-CoA to yield trans-3-enoyl-CoA. Has activity towards short and medium chain 2,4-dienoyl-CoAs, but also towards 2,4,7,10,13,16,19-docosaheptaenoyl-CoA, suggesting that it does not constitute a rate limiting step in the peroxisomal degradation of docosaheptaenoic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. 2,4-dienoyl-CoA reductase subfamily.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

