





LPAAT- γ Monoclonal Antibody

Catalog No	YP-mAb-02658
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	AGPAT3
Protein Name	1-acyl-sn-glycerol-3-phosphate acyltransferase gamma
Immunogen	The antiserum was produced against synthesized peptide derived from human AGPAT3. AA range:121-170
Specificity	LPAAT- γ Monoclonal Antibody detects endogenous levels of LPAAT- γ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AGPAT3; 1-acyl-sn-glycerol-3-phosphate acyltransferase gamma; 1-acylglycerol-3-phosphate O-acyltransferase 3; 1-AGP acyltransferase 3; 1-AGPAT 3; Lysophosphatidic acid acyltransferase gamma; LPAAT-gamma
Observed Band	40kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Nucleus envelope .
Tissue Specificity	Widely expressed with highest levels in testis, pancreas and kidney, followed by spleen, lung, adipose tissue and liver.
Function	catalytic activity:Acyl-CoA + 1-acyl-sn-glycerol 3-phosphate = CoA + 1,2-diacyl-sn-glycerol 3-phosphate.,domain:The HXXXXD motif is essential for acyltransferase activity and may constitute the binding site for the phosphate moiety of the glycerol-3-phosphate.,function:Converts lysophosphatidic acid (LPA) into phosphatidic acid by incorporating an acyl moiety at the sn-2 position of the glycerol backbone.,pathway:Phospholipid metabolism; CDP-diacylglycerol biosynthesis; CDP-diacylglycerol from sn-glycerol 3-phosphate: step 2/3.,similarity:Belongs to the 1-acyl-sn-glycerol-3-phosphate acyltransferase family.,
Background	The protein encoded by this gene is an acyltransferase that converts lysophosphatidic acid into phosphatidic acid, which is the second step in the de



UpingBio technology Co.,Ltd







novo phospholipid biosynthetic pathway. The encoded protein may be an integral membrane protein. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Monoclonal Antibody

Western Blot analysis of various cells using LPAAT- γ

Products Images



