



SOD-2 Monoclonal Antibody

Catalog No	YP-mAb-02883
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SOD2
Protein Name	Superoxide dismutase [Mn] mitochondrial
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human SOD2. AA range:91-140
Specificity	SOD-2 Monoclonal Antibody detects endogenous levels of SOD-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SOD2; Superoxide dismutase [Mn], mitochondrial
Observed Band	24kD
Cell Pathway	Mitochondrion matrix.
Tissue Specificity	Brain,Colon,Heart,Liver,Lung,Mammary carcinoma,Tongue,
Function	catalytic activity:2 superoxide + 2 H(+) = O(2) + H(2)O(2).,cofactor:Binds 1 manganese ion per subunit.,disease:Genetic variation in SOD2 is associated with susceptibility to diabetic nephropathy [MIM:612634]; also called susceptibility to microvascular complications of diabetes type 6 (MVCD6). Diabetic nephropathy is a kidney disease and resultant kidney function impairment due to the long standing effects of diabetes on the microvasculature (glomerulus) of the kidney. Features include increased urine protein and declining kidney function.,function:Destroys radicals which are normally produced within the cells and which are toxic to biological systems.,online information:Superoxide dismutase entry,online information:The Singapore human mutation and polymorphism database,PTM:Nitrated under oxidative stress. Nitration coupled with oxidation inhibits the catalytic activity.,similarity:Belo
Background	This gene is a member of the iron/manganese superoxide dismutase family. It encodes a mitochondrial protein that forms a homotetramer and binds one



manganese ion per subunit. This protein binds to the superoxide byproducts of oxidative phosphorylation and converts them to hydrogen peroxide and diatomic oxygen. Mutations in this gene have been associated with idiopathic cardiomyopathy (IDC), premature aging, sporadic motor neuron disease, and cancer. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been identified on chromosome 1. [provided by RefSeq, Apr 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using SOD-2 Monoclonal Antibody