

## ARRDC2 Monoclonal Antibody


Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse, lgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Catalog No	YP-mAb-03719
Applications       WB         Gene Name       ARRDC2         Protein Name       Arrestin domain-containing protein 2         Immunogen       The antiserum was produced against synthesized peptide derived from human ARRDC2. AA range:141-190         Specificity       ARRDC2 Monoclonal Antibody detects endogenous levels of ARRDC2 protein.         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse,lgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Isotype	IgG
Gene Name       ARRDC2         Protein Name       Arrestin domain-containing protein 2         Immunogen       The antiserum was produced against synthesized peptide derived from human ARRDC2. AA range:141-190         Specificity       ARRDC2 Monoclonal Antibody detects endogenous levels of ARRDC2 protein.         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse,lgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Reactivity	Human;Mouse
Protein Name       Arrestin domain-containing protein 2         Immunogen       The antiserum was produced against synthesized peptide derived from human ARRDC2. AA range:141-190         Specificity       ARRDC2 Monoclonal Antibody detects endogenous levels of ARRDC2 protein.         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse,lgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Applications	WB
Immunogen       The antiserum was produced against synthesized peptide derived from human ARRDC2. AA range:141-190         Specificity       ARRDC2 Monoclonal Antibody detects endogenous levels of ARRDC2 protein.         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse,lgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Gene Name	ARRDC2
ARRDC2. AA range:141-190  Specificity ARRDC2 Monoclonal Antibody detects endogenous levels of ARRDC2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, lgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year	Protein Name	Arrestin domain-containing protein 2
Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Immunogen	
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Specificity	ARRDC2 Monoclonal Antibody detects endogenous levels of ARRDC2 protein.
Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year	Purification	·
Purity ≥90% Storage Stability -20°C/1 year	Dilution	WB 1:500-1:2000
Storage Stability -20°C/1 year	Concentration	1 mg/ml
· · · · · · · · · · · · · · · · · · ·	Purity	≥90%
	Storage Stability	-20°C/1 year
Synonyms ARRDC2; PP2703; Arrestin domain-containing protein 2	Synonyms	ARRDC2; PP2703; Arrestin domain-containing protein 2
Observed Band 44kD	Observed Band	44kD
Cell Pathway plasma membrane,cytoplasmic vesicle,	Cell Pathway	plasma membrane,cytoplasmic vesicle,
Tissue Specificity Brain, Synovial cell,	Tissue Specificity	Brain,Synovial cell,
Function similarity:Belongs to the arrestin family.,	Function	similarity:Belongs to the arrestin family.,
Background similarity:Belongs to the arrestin family.,	Background	similarity:Belongs to the arrestin family.,
matters needing Avoid repeated freezing and thawing! attention	matters needing attention	Avoid repeated freezing and thawing!
<b>Usage suggestions</b> This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.







## **Products Images**



Western Blot analysis of various cells using ARRDC2 Monoclonal Antibody