



# CEP170 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-03758
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	CEP170
<b>Protein Name</b>	Centrosomal protein of 170 kDa
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human CEP170. AA range:701-750
<b>Specificity</b>	CEP170 Monoclonal Antibody detects endogenous levels of CEP170 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CEP170; FAM68A; KAB; KIAA0470; Centrosomal protein of 170 kDa; Cep170; KARP-1-binding protein; KARP1-binding protein
<b>Observed Band</b>	160kD
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole . Cytoplasm, cytoskeleton, spindle . Associated with the mature mother centriole. Associated with spindle microtubules during mitosis. Localizes to the distal appendage region of the centriole (PubMed:31789463). Localizes at the centriole proximal ends (PubMed:31789463). .
<b>Tissue Specificity</b>	Brain,Cervix carcinoma,Epithelium,Skin,Testis,
<b>Function</b>	function:Plays a role in microtubule organization.,PTM:Phosphorylated; probably by PLK1. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the CEP170 family.,similarity:Contains 1 FHA domain.,subcellular location:Associated with the mature mother centriole. Associated with spindle microtubules during mitosis.,subunit:Interacts with PLK1.,
<b>Background</b>	centrosomal protein 170(CEP170) Homo sapiens The product of this gene is a component of the centrosome, a non-membraneous organelle that functions as the major microtubule-organizing center in animal cells. During interphase, the encoded protein localizes to the sub-distal appendages of mature centrioles,



which are microtubule-based structures thought to help organize centrosomes. During mitosis, the protein associates with spindle microtubules near the centrosomes. The protein interacts with and is phosphorylated by polo-like kinase 1, and functions in maintaining microtubule organization and cell morphology. The human genome contains a putative transcribed pseudogene. Several alternatively spliced transcript variants of this gene have been found, but the full-length nature of some of these variants has not been determined. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western Blot analysis of various cells using CEP170 Monoclonal Antibody