



DZIP3 Monoclonal Antibody

Catalog No	YP-mAb-03832
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	DZIP3
Protein Name	E3 ubiquitin-protein ligase DZIP3
Immunogen	The antiserum was produced against synthesized peptide derived from human DZIP3. AA range:681-730
Specificity	DZIP3 Monoclonal Antibody detects endogenous levels of DZIP3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DZIP3; KIAA0675; E3 ubiquitin-protein ligase DZIP3; DAZ-interacting protein 3; RNA-binding ubiquitin ligase of 138 kDa; hRUL138
Observed Band	140kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Widely expressed at low level. Highly expressed in skeletal muscle, kidney and heart. Expressed at low level in placenta, lung, brain, liver and pancreas.
Function	function:E3 Ubiquitin ligase proteins mediate ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Able to specifically bind RNA.,pathway:Protein modification; protein ubiquitination.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Contains 1 RING-type zinc finger.,subunit:Interacts with DAZ proteins.,tissue specificity:Widely expressed at low level. Highly expressed in skeletal muscle, kidney and heart. Expressed at low level in placenta, lung, brain, liver and pancreas.,
Background	function:E3 Ubiquitin ligase proteins mediate ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then



directly transfers the ubiquitin to targeted substrates. Able to specifically bind RNA.,pathway:Protein modification; protein ubiquitination.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Contains 1 RING-type zinc finger.,subunit:Interacts with DAZ proteins.,tissue specificity:Widely expressed at low level. Highly expressed in skeletal muscle, kidney and heart. Expressed at low level in placenta, lung, brain, liver and pancreas.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DZIP3 Monoclonal Antibody