



eIF3 α Monoclonal Antibody

Catalog No	YP-mAb-03844
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	EIF3J
Protein Name	Eukaryotic translation initiation factor 3 subunit J
Immunogen	Synthesized peptide derived from eIF3 α . at AA range: 40-120
Specificity	eIF3 α Monoclonal Antibody detects endogenous levels of eIF3 α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EIF3J; EIF3S1; Eukaryotic translation initiation factor 3 subunit J; eIF3j; Eukaryotic translation initiation factor 3 subunit 1; eIF-3-alpha; eIF3 p35
Observed Band	30kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Epithelium,Fetal liver,Lung carcinoma,Ovarian carci
Function	function:Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. This subunit binds directly within the mRNA entry channel of the 40S ribosome to the aminoacyl (A) site. It may regulate the interaction between the 43S PIC and mRNA.,mass spectrometry: PubMed:17322308,mass spectrometry: PubMed:18599441,P
Background	This gene encodes a core subunit of the eukaryotic initiation factor 3 complex, which participates in the initiation of translation by aiding in the recruitment of



protein and mRNA components to the 40S ribosome. There are pseudogenes for this gene on chromosomes 1, 3, and 9. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2013],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images

SH-SY5Y 293T 3T3

Western Blot analysis of various cells using eIF3 α Monoclonal Antibody



Tubulin 55KD



eIF3 α 30KD