

## FGF-BP2 Monoclonal Antibody

| Catalog No                | YP-mAb-03871  |
|---------------------------|---|
| Isotype                   | IgG   |
| Reactivity                | Human;Rat;Mouse;  |
| Applications              | WB  |
| Gene Name                 | FGFBP2  |
| Protein Name              | Fibroblast growth factor-binding protein 2  |
| Immunogen                 | The antiserum was produced against synthesized peptide derived from human FGFP2. AA range:170-219   |
| Specificity               | FGF-BP2 Monoclonal Antibody detects endogenous levels of FGF-BP2 protein.   |
| Formulation               | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source                    | Monoclonal, Mouse,IgG   |
| Purification              | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution                  | WB 1:500-1:2000   |
| Concentration             | 1 mg/ml   |
| Purity                    | ≥90%  |
| Storage Stability         | -20°C/1 year  |
| Synonyms                  | FGFBP2; KSP37; Fibroblast growth factor-binding protein 2; FGF-BP2; FGF-binding protein 2; FGFBP-2; 37 kDa killer-specific secretory protein; Ksp37; HBp17-related protein; HBp17-RP  |
| Observed Band             | 37kD  |
| Cell Pathway              | Secreted, extracellular space .   |
| Tissue Specificity        | Expressed in serum, peripheral leukocytes and cytotoxic T-lymphocytes, but not in granulocytes and monocytes (at protein level).  |
| Function                  | similarity:Belongs to the fibroblast growth factor-binding protein family.,tissue specificity:Expressed in serum, peripheral leukocytes and cytotoxic T lymphocytes, but not in granulocytes and monocytes (at protein level).,   |
| Background                | This gene encodes a member of the fibroblast growth factor binding protein family. The encoded protein is a serum protein that is selectively secreted by cytotoxic lymphocytes and may be involved in cytotoxic lymphocyte-mediated immunity. An increase in the amount of gene product may be associated with atopic asthma and mild extrinsic asthma.[provided by RefSeq Staff, Oct 2008], |
| matters needing attention | Avoid repeated freezing and thawing!  |
|                           |   |



## UpingBio technology Co.,Ltd








**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using FGF-BP2 Monoclonal Antibody