



# MIPP Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-03960
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	MINPP1
<b>Protein Name</b>	Multiple inositol polyphosphate phosphatase 1
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human MINPP1. AA range:328-377
<b>Specificity</b>	MIPP Monoclonal Antibody detects endogenous levels of MIPP protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	MINPP1; MIPP; Multiple inositol polyphosphate phosphatase 1; 2; 3-bisphosphoglycerate 3-phosphatase; 2,3-BPG phosphatase; Inositol; 1,3,4,5)-tetrakisphosphate 3-phosphatase; Ins(1,3,4,5)P(4) 3-phosphatase
<b>Observed Band</b>	60kD
<b>Cell Pathway</b>	Endoplasmic reticulum lumen .
<b>Tissue Specificity</b>	Widely expressed with highest levels in kidney, liver and placenta.
<b>Function</b>	catalytic activity:Myo-inositol hexakisphosphate + H(2)O = myo-inositol pentakisphosphate (mixed isomers) + phosphate.,disease:Defects in MINPP1 may be involved in follicular thyroid tumors development.,function:Acts as a phosphoinositide 5- and phosphoinositide 6-phosphatase and regulates cellular levels of inositol pentakisphosphate (InsP5) and inositol hexakisphosphate (InsP6) (By similarity). May play a role in bone development (endochondral ossification).,tissue specificity:Widely expressed with highest levels in kidney, liver and placenta.,
<b>Background</b>	This gene encodes multiple inositol polyphosphate phosphatase; an enzyme that removes 3-phosphate from inositol phosphate substrates. It is the only enzyme known to hydrolzye inositol pentakisphosphate and inositol hexakisphosphate. This enzyme also converts 2,3 bisphosphoglycerate (2,3-BPG) to



2-phosphoglycerate; an activity formerly thought to be exclusive to 2,3-BPG synthase/2-phosphatase (BPGM) in the Rapoport-Luebering shunt of the glycolytic pathway.[provided by RefSeq, Sep 2009],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using MIPP Monoclonal Antibody