





MRP-L50 Monoclonal Antibody

IgG Human;Rat;Mouse; WB MRPL50 39S ribosomal protein L50 mitochondrial The antiserum was produced against synthesized peptide derived from human
Human;Rat;Mouse; WB MRPL50 39S ribosomal protein L50 mitochondrial
WB MRPL50 39S ribosomal protein L50 mitochondrial
MRPL50 39S ribosomal protein L50 mitochondrial
39S ribosomal protein L50 mitochondrial
·
The antiserum was produced against synthesized peptide derived from human
MRPL50. AA range:109-158
MRP-L50 Monoclonal Antibody detects endogenous levels of MRP-L50 protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
MRPL50; 39S ribosomal protein L50; mitochondrial; L50mt; MRP-L50
22kD
Mitochondrion .
Carcinoma,Epithelium,Hepatoma,Skin,
Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a putative 39S subunit



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using MRP-L50 Monoclonal Antibody