






PHAPI2 Monoclonal Antibody

Catalog No	YP-mAb-04077
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ANP32B
Protein Name	Acidic leucine-rich nuclear phosphoprotein 32 family member B
Immunogen	The antiserum was produced against synthesized peptide derived from human ANP32B. AA range:1-50
Specificity	PHAPI2 Monoclonal Antibody detects endogenous levels of PHAPI2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ANP32B; APRIL; PHAPI2; Acidic leucine-rich nuclear phosphoprotein 32 family member B; Acidic protein rich in leucines; Putative HLA-DR-associated protein I-2; PHAPI2; Silver-stainable protein SSP29
Observed Band	29kD
Cell Pathway	[Isoform 1]: Nucleus . Cytoplasm . Accumulates in the nuclei at the S phase; [Isoform 2]: Cytoplasm. Lacks a nuclear localization signal.
Tissue Specificity	Expressed in heart, lung, pancreas, prostate and in spleen, thymus and placenta.
Function	function:Multifunctional protein working as a cell cycle progression factor as well as a cell survival factor. Required for the progression from the G1 to the S phase. Anti-apoptotic protein which functions as a caspase-3 inhibitor. Has no phosphatase 2A (PP2A) inhibitor activity.,similarity:Belongs to the ANP32 family.,similarity:Contains 4 LRR (leucine-rich) repeats.,subcellular location:Accumulates in the nuclei at the S phase.,tissue specificity:Expressed in heart, lung, pancreas, prostate and in spleen, thymus and placenta.,
Background	function:Multifunctional protein working as a cell cycle progression factor as well as a cell survival factor. Required for the progression from the G1 to the S phase. Anti-apoptotic protein which functions as a caspase-3 inhibitor. Has no phosphatase 2A (PP2A) inhibitor activity.,similarity:Belongs to the ANP32 family.,similarity:Contains 4 LRR (leucine-rich) repeats.,subcellular



UpingBio technology Co.,Ltd







location: Accumulates in the nuclei at the S phase, tissue specificity: Expressed in heart, lung, pancreas, prostate and in spleen, thymus and placenta.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PHAPI2 Monoclonal Antibody