

## **TIGD1** Monoclonal Antibody

Catalog No	YP-mAb-04242
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TIGD1
Protein Name	Tigger transposable element-derived protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human TIGD1. AA range:251-300
Specificity	TIGD1 Monoclonal Antibody detects endogenous levels of TIGD1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year TIGD1; Tigger transposable element-derived protein 1
	·
Synonyms	TIGD1; Tigger transposable element-derived protein 1
Synonyms Observed Band	TIGD1; Tigger transposable element-derived protein 1 67kD
Synonyms Observed Band Cell Pathway	TIGD1; Tigger transposable element-derived protein 1  67kD  Nucleus .  Brain, Teratocarcinoma,  similarity: Belongs to the tigger transposable element derived protein
Synonyms Observed Band Cell Pathway Tissue Specificity	TIGD1; Tigger transposable element-derived protein 1  67kD  Nucleus .  Brain, Teratocarcinoma,  similarity: Belongs to the tigger transposable element derived protein family., similarity: Contains 1 DDE domain., similarity: Contains 1 HTH CENPB-type DNA-binding domain., similarity: Contains 1 HTH psq-type DNA-binding domain., The protein encoded by this gene belongs to the tigger subfamily of the pogo
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function	TIGD1; Tigger transposable element-derived protein 1  67kD  Nucleus .  Brain, Teratocarcinoma,  similarity: Belongs to the tigger transposable element derived protein family., similarity: Contains 1 DDE domain., similarity: Contains 1 HTH CENPB-type DNA-binding domain., similarity: Contains 1 HTH psq-type DNA-binding domain.,  The protein encoded by this gene belongs to the tigger subfamily of the pogo superfamily of DNA-mediated transposons in humans. These proteins are related to DNA transposons found in fungi and nematodes, and more distantly to the Tc1 and mariner transposases. They are also very similar to the major mammalian centromere protein B. The exact function of this gene is not known. [provided by



## UpingBio technology Co.,Ltd








**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using TIGD1 Monoclonal Antibody