



# ZnT-9 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-04281
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	SLC30A9
<b>Protein Name</b>	Zinc transporter 9
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human SLC30A9. AA range:181-230
<b>Specificity</b>	ZnT-9 Monoclonal Antibody detects endogenous levels of ZnT-9 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	SLC30A9; C4orf1; HUEL; Zinc transporter 9; ZnT-9; Human embryonic lung protein; HuEL; Solute carrier family 30 member 9
<b>Observed Band</b>	63kD
<b>Cell Pathway</b>	Membrane ; Multi-pass membrane protein . Nucleus . Cytoplasm . Cytoplasmic vesicle . Endoplasmic reticulum . Mainly in the cytoplasm (PubMed:10409434). Partial co-localization with endoplasmic reticulum (PubMed:28334855). .
<b>Tissue Specificity</b>	Ubiquitously expressed in fetal and adult tissues and cancer cell lines.
<b>Function</b>	function:Plays a role in the p160 coactivator signaling pathway that mediates transcriptional activation by nuclear receptors (By similarity). Plays a role in transcriptional activation of Wnt-responsive genes.,similarity:Belongs to the cation diffusion facilitator (CDF) transporter (TC 2.A.4) family. SLC30A subfamily.,subunit:Interacts with GRIP1, ESR1 and AR.,tissue specificity:Ubiquitously expressed in fetal and adult tissues and cancer cell lines.,
<b>Background</b>	function:Plays a role in the p160 coactivator signaling pathway that mediates transcriptional activation by nuclear receptors (By similarity). Plays a role in transcriptional activation of Wnt-responsive genes.,similarity:Belongs to the cation diffusion facilitator (CDF) transporter (TC 2.A.4) family. SLC30A subfamily.,subunit:Interacts with GRIP1, ESR1 and AR.,tissue specificity:Ubiquitously expressed in fetal and adult tissues and cancer cell lines.,




**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using ZnT-9 Monoclonal Antibody