



SS-56 Monoclonal Antibody

Catalog No	YP-mAb-04301
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TRIM68
Protein Name	E3 ubiquitin-protein ligase TRIM68
Immunogen	Synthesized peptide derived from the Internal region of human SS-56.
Specificity	SS-56 Monoclonal Antibody detects endogenous levels of SS-56 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TRIM68; GC109; RNF137; SS56; E3 ubiquitin-protein ligase TRIM68; RING finger protein 137; SSA protein SS-56; SS-56; Tripartite motif-containing protein 68
Observed Band	56kD
Cell Pathway	Cytoplasm, perinuclear region. Nucleus. Colocalized with AR in nucleus.
Tissue Specificity	Widely expressed. Expressed at high levels in prostate cancer cell lines. Up-regulation could be restricted to androgen-dependent cells.
Function	disease:Target of immune response in patients with systemic lupus erythematosus (SLE), primary Sjogren syndrome or HIV-1 infection.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,tissue specificity:Detected in all tissues. High expression is noted in spleen and in fetal liver.,
Background	This gene encodes a member of the tripartite motif-containing protein family, whose members are characterized by a "really interesting new gene" (RING) finger domain, a zinc-binding B-box motif, and a coiled-coil region. Members of this family function as E3 ubiquitin ligases and are involved in a broad range of biological processes. This gene regulates the activation of nuclear receptors, such as androgen receptor, and has been implicated in development of prostate cancer cells, where its expression increases in response to a



downregulation of microRNAs. In addition, this gene participates in viral defense regulation as a negative regulator of interferon-beta. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using SS-56 Monoclonal Antibody