



SYT2 Monoclonal Antibody

Catalog No	YP-mAb-05169
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SYT2
Protein Name	Synaptotagmin-2 (Synaptotagmin II) (SytlI)
Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Specificity	SYT2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	46kD
Cell Pathway	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Single-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, chromaffin granule membrane ; Single-pass membrane protein .
Tissue Specificity	Expressed at the neuromuscular junction (PubMed:33659639). Expressed in melanocytes (PubMed:23999003).
Function	cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domains.,domain: The first C2 domain mediates Ca(2+)-dependent phospholipid binding.,domain: The second C2 domain mediates interaction with Stonin 2.,function: May have a regulatory role in the membrane interactions during trafficking of synaptic vesicles at the active zone of the synapse. It binds acidic phospholipids with a specificity that requires the presence of both an acidic head group and a diacyl backbone.,similarity: Belongs to the synaptotagmin family.,similarity: Contains 2 C2 domains.,subcellular location: Synaptic vesicles and chromaffin granules.,subunit: Homotetramer (Probable). Interacts with stonin 2 (By similarity). Interacts with SCAMP5.,
Background	This gene encodes a synaptic vesicle membrane protein. The encoded protein is thought to function as a calcium sensor in vesicular trafficking and exocytosis. Mutations in this gene are associated with myasthenic syndrome, presynaptic, congenital, with or without motor neuropathy. Alternative splicing results in




multiple transcript variants. [provided by RefSeq, Oct 2014],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using SYT2 Monoclonal Antibody