




DHRS7 Monoclonal Antibody

Catalog No	YP-mAb-05512
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	DHRS7 DHRS7A RETSDR4 CGI-86 UNQ285/PRO3448
Protein Name	Dehydrogenase/reductase SDR family member 7 (EC 1.1.-.-) (Retinal short-chain dehydrogenase/reductase 4) (retSDR4)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DHRS7 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	37kD
Cell Pathway	Endoplasmic reticulum membrane . Bound to the endoplasmic reticulum membrane, possibly through a N-terminus anchor. The main bulk of the polypeptide chain was first reported to be facing toward the lumen of the endoplasmic reticulum (PubMed:24246760). However, it was later shown to be facing the cytosol (PubMed:28457967). .
Tissue Specificity	Found predominantly in the adrenal glands, liver, thyroid, prostate, small intestine, colon, stomach, kidney and brain (PubMed:26466768). Lower levels observed in skeletal muscle, the lung and the spleen (PubMed:26466768).
Function	similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,
Background	This gene encodes a member of the short-chain dehydrogenases/reductases (SDR) family, which has over 46,000 members. Members in this family are enzymes that metabolize many different compounds, such as steroid hormones, prostaglandins, retinoids, lipids and xenobiotics. [provided by RefSeq, Apr 2016],
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using DHRS7 Monoclonal Antibody