


NTHL1 Monoclonal Antibody

Catalog No	YP-mAb-05572
Catalog No	
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	NTHL1 NTH1 OCTS3
Protein Name	Endonuclease III-like protein 1 (EC 4.2.99.18)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	NTHL1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	34kD
Cell Pathway	Nucleus . Mitochondrion .
Tissue Specificity	Widely expressed with highest levels in heart and lowest levels in lung and liver.
Function	catalytic activity:The C-O-P bond 3' to the apurinic or apyrimidinic site in DNA is broken by a beta-elimination reaction, leaving a 3'-terminal unsaturated sugar and a product with a terminal 5'-phosphate.,caution:It is uncertain whether Met-1, Met-9 or Met-16 is the initiator.,cofactor:Binds 1 4Fe-4S cluster. The cluster is not important for the catalytic activity, but which is probably involved in the proper positioning of the enzyme along the DNA strand.,developmental stage:Expression levels are regulated during the cell cycle with increased levels during early and mid S-phase.,function:Has both an apurinic and/or apyrimidinic endonuclease activity and a DNA N-glycosylase activity. Incises damaged DNA at cytosines, thymines and guanines. Acts on a damaged strand, 5' from the damaged site. Required for the repair of both oxidative DNA damage and spontaneous mutagenic lesions.,similari
Background	The protein encoded by this gene is a DNA N-glycosylase of the endonuclease III family. Like a similar protein in E. coli, the encoded protein has DNA glycosylase activity on DNA substrates containing oxidized pyrimidine residues



UpingBio technology Co.,Ltd








and has apurinic/apyrimidinic lyase activity. [provided by RefSeq, Oct 2008],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NTHL1 Monoclonal Antibody