


HOME1 Monoclonal Antibody

Catalog No	YP-mAb-05672
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	HOMER1 SYN47
Protein Name	Homer protein homolog 1 (Homer-1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HOME1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 38kD Cytoplasm . Cell junction, synapse, postsynaptic density . Cell junction, synapse . Cell projection, dendritic spine . Isoform 1 inhibits surface expression of GRM5
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 38kD Cytoplasm . Cell junction, synapse, postsynaptic density . Cell junction, synapse . Cell projection, dendritic spine . Isoform 1 inhibits surface expression of GRM5 causing it to be retained in the endoplasmic reticulum
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 28kD Cytoplasm . Cell junction, synapse, postsynaptic density . Cell junction, synapse . Cell projection, dendritic spine . Isoform 1 inhibits surface expression of GRM5 causing it to be retained in the endoplasmic reticulum. Brain,Frontal cortex,Pancreas,Stomach mucosa,Synovial cell, domain:The WH1 domain interacts with the PPXXF motif in GRM1, GRM5, RYR1, RYR2, ITPR1, SHANK 1 and SHANK3.,function:Postsynaptic density scaffolding protein. Binds and cross-links cytoplasmic regions of GRM1, GRM5, ITPR1, DNM3, RYR1, RYR2, SHANK1 and SHANK3. By physically linking GRM1 and GRM5 with ER-associated ITPR1 receptors, it aids the coupling of surface receptors to intracellular calcium release. May also couple GRM1 to Pl3 kinase through its interaction with AGAP2. Isoform 1 regulates the trafficking and surface expression of GRM5. Isoform 3 acts as a natural dominant negative, in dynamic competition with constitutively expressed isoform 1 to regulate synaptic metabotropic glutamate function. Isoform 3, may be involved in the structural changes that occur at synapses during long-lasting neuronal plasticity and



UpingBio technology Co.,Ltd







[provided by RefSeq, Jul 2008],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



