

## NDUA7 Monoclonal Antibody

Catalog No	YP-mAb-05796
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	NDUFA7
Protein Name	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7 (Complex I-B14.5a) (CI-B14.5a) (NADH-ubiquinone oxidoreductase subunit B14.5a)
Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Specificity	NDUA7 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	12kD
Cell Pathway	Mitochondrion inner membrane ; Peripheral membrane protein ; Matrix side .
Tissue Specificity	Lung,Pooled,Umbilical cord blood,
Function	function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFA7 subunit family.,subunit:Complex I is composed of 45 different subunits.,
Background	This gene encodes a subunit of NADH:ubiquinone oxidoreductase (complex I), which is a multiprotein complex located in the inner mitochondrial membrane. Complex I functions in the transfer of electrons from NADH to the respiratory chain. [provided by RefSeq, Mar 2011],
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd








**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using NDUA7 Monoclonal Antibody