



NB5R2 Monoclonal Antibody

Catalog No	YP-mAb-05808
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	CYB5R2
Protein Name	NADH-cytochrome b5 reductase 2 (b5R.2) (EC 1.6.2.2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 120-200
Specificity	NB5R2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	30kD
Cell Pathway	nucleus,mitochondrion,endoplasmic reticulum membrane,membrane,integral component of membrane,
Tissue Specificity	Restricted expression.
Function	catalytic activity:NADH + 2 ferricytochrome b5 = NAD(+) + H(+) + 2 ferrocytochrome b5.,cofactor:FAD.,function:NADH-cytochrome b5 reductases are involved in desaturation and elongation of fatty acids, cholesterol biosynthesis, drug metabolism, and, in erythrocyte, methemoglobin reduction (By similarity). Responsible for NADH-dependent lucigenin chemiluminescence in spermatozoa by reducing both lucigenin and 2-[4-iodophenyl]-3-[4-nitrophenyl]-5-[2,4-disulfophenyl]-2H tetrazolium monosodium salt (WST-1).,similarity:Belongs to the flavoprotein pyridine nucleotide cytochrome reductase family.,similarity:Contains 1 FAD-binding FR-type domain.,tissue specificity:Restricted expression.,
Background	The protein encoded by this gene belongs to the flavoprotein pyridine nucleotide cytochrome reductase family of proteins. Cytochrome b-type NAD(P)H oxidoreductases are implicated in many processes including cholesterol biosynthesis, fatty acid desaturation and elongation, and respiratory burst in neutrophils and macrophages. Cytochrome b5 reductases have soluble and




membrane-bound forms that are the product of alternative splicing. In animal cells, the membrane-bound form binds to the endoplasmic reticulum, where it is a member of a fatty acid desaturation complex. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2014],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using NB5R2 Monoclonal Antibody